

Complexity Theory

Exercise 7: Diagonalisation and Alternation

December 12, 2018

Exercise 7.1. Show that Cook-reducibility is transitive. In other words, show that if **A** is Cook-reducible to **B** and **B** is Cook-reducible to **C**, then **A** is Cook-reducible to **C**.

Exercise 7.2. Show that there exists an oracle **C** such that $\text{NP}^{\text{C}} \neq \text{coNP}^{\text{C}}$.

Hint:

BAKER-CUILL-2010 ASY THEOREM for coNP instead of P .

What kind of Turing machines exist for languages in coNP ? Use the answer to adapt the proof of the

Exercise 7.3. Describe a polynomial-time ATM solving **EXACT INDEPENDENT SET**:

Input: Given a graph G and some number k .

Question: Does there exist a maximal independent set in G of size exactly k ?

Exercise 7.4. Consider the Japanese game *go-moku* that is played by two players **X** and **O** on a 19×19 board. Players alternately place markers on the board, and the first one to have five of its markers in a row, column, or diagonal wins.

Consider the generalised version of *go-moku* on an $n \times n$ board. Say that a *position* of *go-moku* is a placement of markers on such a board as it could occur during the game. Define

$$\mathbf{GM} = \{ \langle B \rangle \mid B \text{ is a position of go-moku where X has a winning strategy} \}.$$

Describe a polynomial-time ATM solving **GM**.

Exercise 7.5. Show that $\text{AEXP TIME} = \text{EXPSPACE}$.