

Hannes Strass

Faculty of Computer Science, Institute of Artificial Intelligence, Computational Logic Group

Cooperative Games: Definition and the Core

Lecture 11, 1st Jul 2024 // Algorithmic Game Theory, SS 2024

Previously ...

- **General Game Playing** is concerned with computers learning to play previously unknown games without human intervention.
- The **game description language** (GDL) is used to declaratively specify (deterministic) games (with complete information about game states).
- The syntax of GDL game descriptions is that of **normal logic programs**; various restrictions apply to obtain a finite, unique interpretation.
- The semantics of GDL is given through a state transition system.
- GDL-II allows to represent moves by **Nature** and information sets.
- The semantics of GDL-II can be given through extensive-form games.
- Conversely, GDL-II can express any finite extensive-form game.

Written Exam

16th Aug 2024, 13:00–14:30 HSZ/E01/U

Overview

Cooperative Games with Transferable Utility

Solution Concept: The Core
 ϵ -Cores and the Least Core
The Cost of Stability

Cooperative Games: Motivation

- In a **noncooperative game**, players cannot enter binding agreements.
- (Players can still cooperate if it pays off for them.)
- In a **cooperative game**, players form explicit **coalitions**.
- The coalition gets some (overall) payoff, which is then to be distributed among the coalition's members (transferable utility).
- Players are still assumed to be rationally maximising their **individual payoffs**.

Example: Hospitals and X-Ray Machines

- Three hospitals (in the same city) are planning to buy x-ray machines.
- However, not every hospital necessarily needs its own machine.
- The smallest machine costs $\$5m$ and could cover the needs of any two hospitals.
- A larger machine costs $\$9m$ and could cover the needs of all three hospitals.
- Hospitals forming a coalition C can jointly save the difference to each individual hospital $i \in C$ buying its own $\$5m$ machine.
- It is in society's interest to save money while covering patients' needs.

What should the hospitals do?

Cooperative Games with Transferable Utility

Cooperative Games with Transferable Utility

Definition

A **cooperative game with transferable utility** is a pair $G = (P, v)$ where

- $P = \{1, 2, \dots, n\}$ is the set of players and
 - $v: 2^P \rightarrow \mathbb{R}_{\geq 0}$ is the **characteristic function** of G .
-
- **Intuition:** Coalition $C \subseteq P$ earns $v(C)$ by cooperating.
 - **Terminology:** We will occasionally omit “with transferable utility”.

Assumption

For any cooperative game $G = (P, v)$, we have:

1. Normalisation: $v(\emptyset) = 0$.
2. Monotonicity: $C \subseteq D \subseteq P$ implies $v(C) \leq v(D)$.

Note that a cooperative game with n players requires a representation of a size that is exponential in n .

Cooperative Games: Example

Hospitals and X-Ray Machines

Three hospitals are planning to buy x-ray machines. However, not every hospital necessarily needs its own machine. A small machine costs \$5m and could cover the needs of any two hospitals. A larger machine costs \$9m and could cover the needs of all three hospitals. Hospitals forming a coalition C can jointly save the difference to each individual hospital $i \in C$ buying its own \$5m machine.

- $P = \{1, 2, 3\}$,
- $v(P) = 6$,
- $v(C) = 5$ for $|C| = 2$,
- $v(\{i\}) = 0$ for $i \in P$.

Coalition Structure

Definition

Let $G = (P, v)$ be a cooperative game (with transferable utility).

A **coalition structure** for G is a partition $\mathcal{C} = \{C_1, \dots, C_k\}$ of P , that is,

- $C_1, \dots, C_k \subseteq P$,
 - $C_1 \cup \dots \cup C_k = P$, and
 - $C_i \cap C_j = \emptyset$ for all $1 \leq i \neq j \leq k$.
-
- The coalition structure $\mathcal{C} = \{P\}$ is called the **grand coalition**.
 - $v(C)$ is the **collective payoff** of a coalition; it remains to be specified how to distribute the gains to the coalition's members.

Hospitals and X-Ray Machines

For $P = \{1, 2, 3\}$, some possible coalition structures are $\mathcal{C}_1 = \{\{1, 2, 3\}\}$, $\mathcal{C}_2 = \{\{1, 3\}, \{2\}\}$, and $\mathcal{C}_3 = \{\{1\}, \{2\}, \{3\}\}$.

Outcome of a Cooperative Game

Definition

Let $G = (P, v)$ be a cooperative game (with transferable utility).

An **outcome** of $G = (P, v)$ is a pair $(\mathcal{C}, \mathbf{a})$ where \mathcal{C} is a coalition structure and $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}^n$ is a payoff vector such that $a_i \geq 0$ for each $i \in P$ and

$$\sum_{i \in C} a_i = v(C) \quad \text{for each coalition } C \in \mathcal{C}.$$

Efficiency: For each coalition $C \in \mathcal{C}$, its payoff $v(C)$ is distributed completely.

Transferable Utility: Players within coalitions can transfer payoffs freely.

Hospitals and X-Ray Machines: Outcomes

$\mathcal{C}_1 = \{\{1, 2, 3\}\}$ with $\mathbf{a}_1 = (2, 2, 2)$, $\mathcal{C}_2 = \{\{1, 3\}, \{2\}\}$ with $\mathbf{a}_2 = (2.5, 0, 2.5)$,
 $\mathcal{C}_3 = \{\{1\}, \{2\}, \{3\}\}$ with $\mathbf{a}_3 = (0, 0, 0)$, but also \mathcal{C}_2 with $\mathbf{a}'_2 = (3, 0, 2)$.

No outcome: \mathcal{C}_2 with $(2, 1, 2)$.

Superadditive Games (1)

Definition

Let $G = (P, v)$ be a cooperative game (with transferable utility).

G is called **superadditive** iff for all coalitions $C, D \subseteq P$

$$C \cap D = \emptyset \quad \text{implies} \quad v(C \cup D) \geq v(C) + v(D).$$

Intuition: $C \cup D$ can achieve what C and D can achieve separately; there might be additional synergistic effects.

Non-Example

- A group C of emacs-using programmers achieves a part of a task T in 8h.
- A (disjoint) group D of vi-using programmers achieves the rest of T in 8h.
- The group $C \cup D$, attempting to work together, might not achieve T in 8h.

We will only consider superadditive games unless specified otherwise.

Superadditive Games (2)

Observation

Let $G = (P, v)$ be a superadditive (cooperative) game.
For every coalition structure $\mathcal{C} = \{C_1, \dots, C_k\}$, we have

$$v(P) \geq v(C_1) + \dots + v(C_k)$$

↪ In superadditive games, we can expect the grand coalition to form.
However, it does not automatically mean that the grand coalition is “stable”:

Example

- The “Hospitals and X-Ray Machines” game is superadditive.
- In outcome $(\{\{1, 2, 3\}\}, (2, 2, 2))$, e.g. $\{1, 2\}$ have an incentive to deviate:
- In $(\{\{1, 2\}, \{3\}\}, (2.5, 2.5, 0))$, they would increase their individual payoff.

↪ It remains to analyse how to distribute the grand coalition’s payoff.

Solution Concept: The Core

Imputations

Definition

Let $G = (P, v)$ be a cooperative game (with transferable utility).

- A payoff vector $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}^n$ is **individually rational** iff

$$a_i \geq v(\{i\}) \quad \text{for all } i \in P.$$

- A payoff vector \mathbf{a} is **efficient** (w.r.t. the grand coalition) iff

$$\sum_{i=1}^n a_i = v(P)$$

- An **imputation** for G is a payoff vector \mathbf{a} that is efficient and individually rational. The set of all imputations of G is denoted $Imp(G)$.


Observations

1. $Imp(G) \neq \emptyset$ iff $v(P) \geq \sum_{i \in P} v(\{i\})$.
2. If G is superadditive, then $Imp(G) \neq \emptyset$.

Imputations: Visualisation

Consider the game $G = (P, v)$ with:

- $P = \{A, B, C\}$,
- $v(P) = 100$ and $v(\{i\}) = 0$ for $i \in P$,
- $v(\{A, B\}) = v(\{A, C\}) = 50$, and $v(\{B, C\}) = 30$.



The Core of a Cooperative Game

Definition

Let $G = (P, v)$ be a cooperative game (with transferable utility).

1. An imputation $(a_1, \dots, a_n) \in \text{Imp}(G)$ is **coalitionally rational** iff

$$\sum_{i \in C} a_i \geq v(C) \text{ for all coalitions } C \subseteq P.$$

2. The **core** of G is the set of all coalitionally rational imputations for G .

Intuition: No group C has an incentive to break off the grand coalition.

Example


In “Hospitals and X-Ray Machines”, the core is empty:

- If $(a_1, a_2, a_3) \in \text{Core}(G)$, then $a_1 + a_2 + a_3 = 6$ by being an imputation.
- But for any $i, j \in \{1, 2, 3\}$ with $i \neq j$ we also have $a_i + a_j \geq v(\{a_i, a_j\}) = 5$.
- Let $a_i \leq a_j \leq a_k$, then $a_i + a_j \geq 5$, but $a_k \leq 1$ and $a_i + a_j \leq 2$, contradiction.

The Core: Visualisation

Consider the game $G = (P, v)$ with:

- $P = \{A, B, C\}$,
- $v(P) = 100$ and $v(\{i\}) = 0$ for $i \in P$,
- $v(\{A, B\}) = v(\{A, C\}) = 50$, and $v(\{B, C\}) = 30$.



Cores of Cooperative Games: Example (1)

Chess Pairings

A group of $n \geq 3$ people want to play chess. Every pair of players appointed to play against each other receives \$1.

$$P = \{1, \dots, n\}$$
$$v(C) = \begin{cases} \frac{|C|}{2} & \text{if } |C| \text{ is even,} \\ \frac{|C|-1}{2} & \text{otherwise} \end{cases}$$

- For $n \geq 4$ even, the payoff vector $\mathbf{a}_n := \left(\frac{1}{2}, \dots, \frac{1}{2}\right)$ is in the core:
 - deviation by an odd group $C \subseteq P$ would yield $v(C) = \frac{|C|-1}{2} < \frac{1}{2} \cdot |C|$;
 - deviation by an even group $C \subseteq P$ would yield $v(C) = \frac{|C|}{2} = \frac{1}{2} \cdot |C|$.
- In fact, for $n \geq 4$ even, we have $\text{Core}(G) = \{\mathbf{a}_n\}$:
 - If $\mathbf{a} \in \text{Core}(G)$, then for any $\{a_i, a_j\} \subseteq P$, it follows that $a_i + a_j \geq v(\{a_i, a_j\}) = 1$.
 - From $\mathbf{a} \in \text{Imp}(G)$, we get $a_1 + \dots + a_n = \frac{n}{2}$, and we obtain $a_i = \frac{1}{2}$ for all $i \in P$.
- For $n \geq 3$ odd, the core is empty: (One player remains without a partner.)
 - For $n = 3$ and $\mathbf{a} \in \text{Core}(G)$, we get $a_1 + a_2 + a_3 = 1$, so e.g. $a_1 > 0$.
 - But then $a_2 + a_3 = 1 - a_1 < 1$ although $v(\{a_2, a_3\}) = 1$, contradicting $\mathbf{a} \in \text{Core}(G)$.

Cores of Cooperative Games: Example (2)

Shoe Makers

Of 201 shoe makers, (the first) 100 have made one left shoe each, (the remaining) 101 have made one right shoe each. A pair of shoes consists of one left and one right shoe (ignoring sizes), and can be sold for \$10.

$$P = \{1, 2, \dots, 201\}$$

$$v(C) = 10 \cdot \min\{|C_L|, |C_R|\}$$

where

$$C_L := \{c \in C \mid c \leq 100\}$$

$$C_R := \{c \in C \mid c \geq 101\}$$

- The grand coalition makes a total of \$1000 from selling all 100 pairs.
- The core of this game contains as only imputation $\mathbf{a} = (a_1, a_2, \dots, a_{201})$ with $a_1 = a_2 = \dots = a_{100} = 10$ and $a_{101} = a_{102} = \dots = a_{201} = 0$:
- For any imputation \mathbf{b} with $b_i > 0$ for some $101 \leq i \leq 201$, the coalition $P \setminus \{i\}$ would obtain $v(P \setminus \{i\}) = v(P) > v(P) - b_i = \sum_{j \in P, j \neq i} b_j$ on their own.
- **Intuitively:** Left shoes are scarce, right shoes are overabundant.

The Core is a Convex Set

Proposition

Let $G = (P, v)$ be a cooperative game (with transferable utility).

For any $\mathbf{a}, \mathbf{b} \in \text{Core}(G)$ and $\alpha \in [0, 1]$, we have $\alpha \cdot \mathbf{a} + (1 - \alpha) \cdot \mathbf{b} \in \text{Core}(G)$.

Proof.

1. It is clear that $\sum_{i \in P} a_i = \sum_{i \in P} b_i = v(P)$, and that

$$\sum_{i \in P} (\alpha \cdot a_i + (1 - \alpha) \cdot b_i) = \alpha \cdot \sum_{i \in P} a_i + (1 - \alpha) \cdot \sum_{i \in P} b_i = \alpha \cdot v(P) + (1 - \alpha) \cdot v(P) = v(P)$$

2. Let $C \subseteq P$. With $\sum_{i \in P} a_i \geq v(C)$ and $\sum_{i \in P} b_i \geq v(C)$ we get

$$\sum_{i \in P} (\alpha \cdot a_i + (1 - \alpha) \cdot b_i) = \alpha \cdot \sum_{i \in P} a_i + (1 - \alpha) \cdot \sum_{i \in P} b_i \geq \alpha \cdot v(C) + (1 - \alpha) \cdot v(C) = v(C)$$

□

Linear Programming (in a Nutshell)

Definition

- A **linear program** is of the form

$$\begin{aligned} & \text{maximise} && \mathbf{c}^T \mathbf{x} \\ & \text{subject to} && \mathbf{Ax} \leq \mathbf{b}, \\ & && \mathbf{x} \geq 0, \\ & && \text{and } \mathbf{x} \in \mathbb{R}^k \end{aligned}$$

where \mathbf{x} is a vector of **decision variables**, and \mathbf{A} , \mathbf{b} , \mathbf{c} are a matrix and two vectors of real values; the expression $\mathbf{c}^T \mathbf{x}$ is the **objective function**.

- If there is no objective function the program is a **feasibility problem**.
- A **solution** is a variable-value assignment that satisfies all constraints.
- A linear program is a special case of a mixed integer program (Lecture 2).
- Linear programming problems can be solved in polynomial time.

Computing the Core

For a given cooperative game $G = (P, v)$, its core is given by the feasible region of the following linear program over variables a_1, \dots, a_n :

$$\begin{array}{ll} \text{find} & a_1, \dots, a_n \\ \text{subject to} & a_j \geq 0 \quad \text{for all } i \in P \\ & \sum_{i \in P} a_i = v(P) \\ & \sum_{i \in C} a_i \geq v(C) \quad \text{for all } C \subseteq P \end{array}$$

Observe: The problem specification contains $2^n + n + 1$ constraints.

Corollary

For a cooperative game $G = (P, v)$ whose characteristic function v is explicitly represented, its core can be computed in deterministic polynomial time.

The ε -Core

Definition

Let $G = (P, v)$ be a cooperative game (with transferable utility) and $\varepsilon \in \mathbb{R}$.

1. The set of **pre-imputations of G** is

$$\text{PreImp}(G) := \{(a_1, \dots, a_n) \in \mathbb{R}^n \mid \sum_{i \in P} a_i = v(P)\}$$

2. The **ε -core of G** is the following set:

$$\varepsilon\text{-Core}(G) := \left\{ (a_1, \dots, a_n) \in \text{PreImp}(G) \mid \sum_{i \in C} a_i \geq v(C) - \varepsilon \text{ for all } C \subseteq P \right\}$$

- **Intuition:** Coalitions $C \subsetneq P$ that leave P have to pay a penalty of at least ε .
- For $\varepsilon = 0$, we have $0\text{-Core}(G) = \text{Core}(G)$.
- If $\text{Core}(G) = \emptyset$, then there is some $\varepsilon \in \mathbb{R}$, $\varepsilon > 0$, for which $\varepsilon\text{-Core}(G) \neq \emptyset$.
- If $\text{Core}(G) \neq \emptyset$, then there is some $\varepsilon \in \mathbb{R}$, $\varepsilon < 0$, for which $\varepsilon\text{-Core}(G) = \emptyset$.

The Least Core

Definition

Let $G = (P, v)$ be a cooperative game (with transferable utility).
The **least core of G** is the intersection of all non-empty ε -cores of G .

Alternatively: The least core of G is $\tilde{\varepsilon}$ -Core(G) for $\tilde{\varepsilon} \in \mathbb{R}$ such that $\tilde{\varepsilon}$ -Core(G) $\neq \emptyset$ and ε -Core(G) = \emptyset for all $\varepsilon < \tilde{\varepsilon}$.

The value of the least core can be computed via linear programming:

$$\begin{array}{ll} \text{minimise} & \varepsilon \\ \text{subject to} & a_i \geq 0 \quad \text{for all } i \in P \\ & \sum_{i \in P} a_i = v(P) \\ & \sum_{i \in C} a_i \geq v(C) - \varepsilon \quad \text{for all } C \subseteq P \end{array}$$

The Cost of Stability

Idea: If $\text{Core}(G) = \emptyset$, stabilise G by subsidising the grand coalition.

Modelling Assumptions

- Some external authority has an interest in a stable grand coalition.
- The supplemental payment γ gets distributed among P along with $v(P)$.

Definition

Let $G = (P, v)$ be a cooperative game (with transferable utility).

1. For a supplemental payment $\gamma \geq 0$, the **adjusted game** $G_\gamma = (P, v')$ has

$$v'(C) := \begin{cases} v(P) + \gamma & \text{if } C = P, \\ v(C) & \text{otherwise.} \end{cases}$$

2. The **cost of stability of G** is $\inf \{ \gamma \in \mathbb{R} \mid \gamma \geq 0 \text{ and } \text{Core}(G_\gamma) \neq \emptyset \}$.

Computing the Cost of Stability

Example: Hospitals and X-Ray Machines

The cost of stability is $\gamma = 1.5$: In G_γ , we have $v'(\{1, 2, 3\}) = 6 + 1.5 = 7.5$, whence for no $C \subseteq \{1, 2, 3\}$ with $|C| = 2$ it would pay to deviate (as $v'(C) = 5$).

The cost of stability can be computed by linear programming:

$$\begin{array}{ll} \text{minimise} & \gamma \\ \text{subject to} & \gamma \geq 0 \\ & a_i \geq 0 \quad \text{for all } i \in P \\ & \sum_{i \in P} a_i = v(P) + \gamma \\ & \sum_{i \in C} a_i \geq v(C) \quad \text{for all } C \subseteq P \end{array}$$

Least Core vs. Cost of Stability

Observation

For any cooperative game G , the following are equivalent:

1. $\text{Core}(G) = \emptyset$.
2. The value ε of the least core is strictly positive.
3. The cost γ of stability is strictly positive.

What is the relationship between the values ε and γ ?

- **Least core:** Punish undesired behaviour
↪ a fine for leaving the grand coalition.
- **Cost of stability:** Encourage desired behaviour
↪ a subsidy for staying in the grand coalition.

Least Core v. Cost of Stability: Examples

Let $n \geq 2$ and consider the following two games (i.e. where $P = \{1, \dots, n\}$):

$$G_1 = (P, v_1)$$

$$v_1(C) = \begin{cases} n-1 & \text{if } C \cap \{1, 2\} \neq \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

$$G_2 = (P, v_2)$$

$$v_2(C) = \begin{cases} 1 & \text{if } C \neq \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

$$G_3 = (P, v_3)$$

$$v_3(C) = \begin{cases} \frac{2n-2}{n} & \text{if } C \cap \{1, 2\} \neq \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

- In both games G_1 and G_2 , the core is empty.
- The cost of stability in both games G_1 and G_2 is $\gamma = n - 1$:
 $\mathbf{a}_1 = (n - 1, n - 1, 0, \dots, 0)$ vs. $\mathbf{a}_2 = (1, 1, 1, \dots, 1)$
- The value of the least core in G_1 is $\varepsilon_1 = \frac{n-1}{2}$, via $\left(\frac{n-1}{2}, \frac{n-1}{2}, 0, \dots, 0\right)$.
- The value of the least core in G_2 is $\varepsilon_2 = \frac{n-1}{n}$, via $\left(\frac{1}{n}, \frac{1}{n}, \dots, \frac{1}{n}\right)$.
- For G_3 , we have $\varepsilon_3 = \frac{n-1}{n}$ via $\mathbf{a}_3 = \left(\frac{n-1}{n}, \frac{n-1}{n}, 0, \dots, 0\right)$ and $\gamma_3 = \frac{2n-2}{n}$ via $\mathbf{a}'_3 = \left(\frac{2n-2}{n}, \frac{2n-2}{n}, 0, \dots, 0\right)$.

Conclusion

Summary

- In **cooperative** games, players P form explicit **coalitions** $C \subseteq P$.
- Coalitions receive payoffs, which are distributed among its members.
- We concentrate on **superadditive** games, where disjoint coalitions can never decrease their payoffs by joining together.
- Of particular interest is the **grand coalition** $\{P\}$ and whether it is *stable*.
- An **imputation** is an outcome that is efficient and individually rational.
- Various solution concepts formalise stability of the grand coalition:
 - the **core** contains all imputations where no coalition has an incentive to leave;
 - the **ε -core** disincentivises leaving the grand coalition via a fine of ε ;
 - the **cost of stability** subsidises staying in the grand coalition via a bonus γ .
- The core is a convex set.